

# Quantum Information Processing Assignment 1

Name: Nate Stemen (20906566)  
 Email: nate.stemen@uwaterloo.ca

Due: Thur, Sep 17, 2020 11:59 PM  
 Course: QIC 710

## Problem 1

Distinguishing between pairs of qubit states.

- (a)  $|0\rangle$  and  $|+\rangle$
- (b)  $|0\rangle$  and  $-\frac{1}{2}|0\rangle + \frac{\sqrt{3}}{2}|1\rangle$
- (c)  $\frac{1}{\sqrt{2}}|0\rangle + \frac{i}{\sqrt{2}}|1\rangle$  and  $\frac{1}{\sqrt{2}}|0\rangle - \frac{i}{\sqrt{2}}|1\rangle$

**Solution.** (a) To best distinguish between these two states we will apply  $R_\theta$  with  $\theta = \frac{\pi}{8}$ . Under this transformation the two states get transformed as follows.

$$R_\theta |0\rangle = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \qquad R_\theta |+\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} \cos \theta - \sin \theta \\ \cos \theta + \sin \theta \end{bmatrix}$$

Under this transformation, we have the following probability table.

	Probability of measuring state	
	0\rangle	1\rangle
$R_\theta  0\rangle$	$\cos^2 \theta \approx 0.85$	$\sin^2 \theta \approx 0.15$
$R_\theta  +\rangle$	$\frac{1}{2}(\cos \theta - \sin \theta)^2 \approx 0.15$	$\frac{1}{2}(\cos \theta + \sin \theta)^2 \approx 0.85$

This leaves us with an overall success probability of 85%.

(b) Here we will make a clockwise rotation of  $15^\circ$  which we will do with  $R_\theta$  with  $\theta = -\frac{\pi}{12}$ . Under this rotation the states are transformed as follows (using  $|\psi\rangle = -\frac{1}{2}|0\rangle + \frac{\sqrt{3}}{2}|1\rangle$  for notational convenience).

$$R_\theta |0\rangle = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \qquad R_\theta |\psi\rangle = \frac{-1}{2} \begin{bmatrix} \cos \theta + \sqrt{3} \sin \theta \\ \sin \theta - \sqrt{3} \cos \theta \end{bmatrix}$$

With these transformed states we can now evaluate the success probabilities given each state as above.

	Probability of measuring state	
	0\rangle	1\rangle
$R_\theta  0\rangle$	$\cos^2 \theta \approx 0.93$	$\sin^2 \theta \approx 0.07$
$R_\theta  \psi\rangle$	$\left  \frac{-1}{4} (\cos \theta + \sqrt{3} \sin \theta)^2 \right  \approx 0.07$	$\left  \frac{-1}{4} (\sin \theta - \sqrt{3} \cos \theta)^2 \right  \approx 0.93$

So here our overall success probability would be 93%. Pretty solid I do say.

(c) These points are anti-podal on the Bloch sphere, and hence orthogonal, which means they are perfectly distinguishable. To find the unitary transformation  $U$  that allows us to measure the state in the computational basis, we need it to satisfy the following two equations.

$$U \left( \frac{1}{\sqrt{2}} |0\rangle + \frac{i}{\sqrt{2}} |1\rangle \right) = |0\rangle$$

$$U \left( \frac{1}{\sqrt{2}} |0\rangle - \frac{i}{\sqrt{2}} |1\rangle \right) = |1\rangle$$

These two equations are equivalent to the following two matrix equations.

$$\frac{1}{\sqrt{2}} \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 \\ i \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$
$$\frac{1}{\sqrt{2}} \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 \\ -i \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

With the matrix written this way we have 4 equations with 4 unknowns. Solving for  $U$  gives us

$$U = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -i \\ 1 & i \end{bmatrix}.$$

This transformation maps our two states to the computational basis which allows us to measure and distinguish perfectly between the two states and hence have 100% success probability.

**Problem 2**

Product states versus entangled states.

$$(a) \frac{1}{2} |00\rangle + \frac{1}{2} |01\rangle + \frac{1}{2} |10\rangle - \frac{1}{2} |11\rangle$$

$$(b) \frac{1}{2} |00\rangle + \frac{1}{2} |01\rangle - \frac{1}{2} |10\rangle - \frac{1}{2} |11\rangle$$

$$(c) \frac{1}{\sqrt{3}} |00\rangle + \frac{1}{\sqrt{3}} |01\rangle + \frac{1}{\sqrt{3}} |10\rangle$$

**Solution.** For this problem we will use the fact that the tensor products of two general single qubit states is  $\alpha_0\beta_0 |00\rangle + \alpha_0\beta_1 |01\rangle + \alpha_1\beta_0 |10\rangle + \alpha_1\beta_1 |11\rangle$ .

(a) With the above formula we have the following four equations.

$$\alpha_0\beta_0 = \frac{1}{2} \tag{1}$$

$$\alpha_0\beta_1 = \frac{1}{2} \tag{2}$$

$$\alpha_1\beta_0 = \frac{1}{2} \tag{3}$$

$$\alpha_1\beta_1 = -\frac{1}{2} \tag{4}$$

Now if we divide eq. (1) by eq. (2) we get  $\beta_0 = \beta_1$  which we will call  $\beta$ . Dividing eq. (3) by eq. (4) yields  $\frac{\alpha_1\beta}{\alpha_1\beta} = \frac{1/2}{-1/2} \implies 1 = -1$ . With this we can conclude this state is not a possible product state and hence an **entangled state**.

(b) The following tensor product gives rise to the desired state which tells us the state is a **product state**, not an entangled state.

$$\left( \frac{1}{\sqrt{2}} |0\rangle - \frac{1}{\sqrt{2}} |1\rangle \right) \otimes \left( \frac{1}{\sqrt{2}} |0\rangle + \frac{1}{\sqrt{2}} |1\rangle \right) = \frac{1}{2} |00\rangle + \frac{1}{2} |01\rangle - \frac{1}{2} |10\rangle - \frac{1}{2} |11\rangle$$

(c) Given the  $|11\rangle$  term is nonexistent, we know  $\alpha_1\beta_1 = 0$ . This implies  $\alpha_1 = 0$  or  $\beta_1 = 0$ . If this was the case then either the  $\alpha_1\beta_0 |10\rangle$  term would be 0 or the  $\alpha_0\beta_1 |01\rangle$  term would be zero. This is not the case so we conclude this is an **entangled state**.