

# Advanced Quantum Theory Homework 1

**Name:** Nate Stemen (20906566)  
**Email:** nate.stemen@uwaterloo.ca

**Due:** Fri, Sep 18, 2020 11:59 AM  
**Course:** AMATH 673

## Exercise 1

Prove  $\{c, f\} = 0$ .

**Solution.** The antisymmetry of the Poisson bracket allows us to equivalently prove  $\{f, c\} = 0$ .

$$\begin{aligned}\{f, c\} &= \{f, \sqrt{c}\}\sqrt{c} + \sqrt{c}\{f, \sqrt{c}\} \\ &= 2\sqrt{c}\{f, \sqrt{c}\} \\ &= -2\sqrt{c}\{\sqrt{c}, f\} \\ &= -2\{c, f\} \\ &= 2\{f, c\}\end{aligned}$$

Since we now have  $\{f, c\} = 2\{f, c\}$  we can subtract  $\{f, c\}$  from both sides to obtain  $\{f, c\} = 0$ .

## Exercise 2

Show that  $\{f, f\} = 0$  for any  $f$ .

**Solution.** Again using the antisymmetry axiom we can see  $\{f, f\} = -\{f, f\}$  and hence adding  $\{f, f\}$  to both sides we obtain  $\{f, f\} = 0$ .

## Exercise 3

Assume that  $n$  is a positive integer.

- (a) Evaluate  $\{x_1, p_1^n\}$
- (b) Evaluate  $\{x_1^n, p_1\}$

**Solution.** ?? I will use a proof by induction to prove  $\{x_1, p_1^n\} = (2p_1)^{n-1}$ . Starting with the base case of  $n = 1$ . Then we have  $\{x_1, p_1\} = (2p_1)^0 = 1$  which agrees with the definition of the Poisson bracket for position and momentum. Assuming the formula is true for  $n$  we will show it's true for  $n + 1$ .

$$\begin{aligned}\{x_1, p_1^{n+1}\} &= \{x_1, p_1^n\}p_1 + p_1\{x_1, p_1^n\} \\ &= 2p_1\{x_1, p_1^n\} \\ &= 2p_1(2p_1)^{n-1} \\ &= (2p_1)^n = (2p_1)^{(n+1)-1}\end{aligned}$$

?? I know this is no longer part of the homework, but I did it before the changes were made, so I figured I would leave it in.

Here we will use the fact that  $\{fg, h\} = \{f, h\}g + f\{g, h\}$  which can be derived from the product rule as follows.

$$\begin{aligned}\{fg, h\} &= -\{h, fg\} \\ &= -\{h, f\}g - f\{h, g\} \\ &= \{f, h\}g + f\{g, h\}\end{aligned}$$

Again I will use proof by induction to prove  $\{x_2^n, p_2\} = nx_2^{n-1}$ . Starting with the base case of  $n = 1$ . Then we have  $\{x_2, p_2\} = 1 \cdot x_2^0 = 1$  which agrees with the definition. Assuming the formula is true for  $n$  we will show it's true for  $n + 1$ .

$$\begin{aligned}\{x_2^{n+1}, p_2\} &= \{x_2^n, p_2\}x_2 + x_2^n\{x_2, p_2\} \\ &= (nx_2^{n-1})x_2 + x_2^n \\ &= nx_2^n + x_2^n \\ &= (n + 1)x_2^n = (n + 1)x_2^{(n+1)-1}\end{aligned}$$

#### Exercise 4

Verify  $\{-2p_1, 3x_1^2 + 7p_3^4 - 2x_2^2p_1^3 + 6\} = 12x_1$ .

**Solution.**

$$\begin{aligned}\text{LHS} &= -2\{p_1, 3x_1^2 + 7p_3^4 - 2x_2^2p_1^3 + 6\} \\ &= -2\left(3\{p_1, x_1^2\} + 7\{p_1, p_3^4\} - 2\{p_1, x_2^2p_1^3\} + \{p_1, 6\}\right) \\ &= -2\left(-6x_1 - 2\left(\{p_1, x_2^2\}p_1^3 + x_2^2\{p_1, p_1^3\}\right)\right) \\ &= 12x_1\end{aligned}$$

#### Exercise 5

Show that the Poisson bracket is not associative by giving a counterexample.

**Solution.** If the Poisson bracket was associative it would mean the following:  $\{f, \{g, h\}\} = \{\{f, g\}, h\}$  for  $f, g, h$  polynomials in  $x_i, p_i$ . Take  $f = p_1^2$ ,  $g = x_1$ , and  $h = p_1$ . We can then evaluate both sides to see this does not hold.

$$\begin{aligned}\text{LHS} &= \{p_1^2, \{x_1, p_1\}\} = \{p_1^2, 1\} = 0 \\ \text{RHS} &= \{\{p_1^2, x_1\}, p_1\} = \{-2p_1, p_1\} = 1\end{aligned}$$

Hence the Poisson bracket is not associative.

**Exercise 6**

Look up and state the axioms of

- a Lie algebra
- an associative algebra
- a Poisson algebra

**Solution.** ?? A *Lie algebra* is a vector space  $\mathfrak{g}$  equipped with bilinear map (called a Lie Bracket)  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  which is anticommutative ( $[x, y] = -[y, x]$ ) and satisfies the Jacobi Identity.

$$[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0 \quad \forall x, y, z \in \mathfrak{g}$$

?? If  $R$  is a commutative ring, then an *associative algebra* is an  $R$ -module  $V$  together with a bilinear map  $V \times V \rightarrow V$  that is associative and has an identity.

?? A *Poisson algebra* is a vector space equipped with two bilinear products  $- \cdot -$  and  $\{\cdot, \cdot\}$  such that

- $- \cdot -$  forms an associative algebra,
- $\{\cdot, \cdot\}$  is antisymmetric, obeys the Jacobi identity and forms a Lie algebra
- The Poisson bracket  $\{\cdot, \cdot\}$  acts as  $\{x, y \cdot z\} = \{x, y\} \cdot z + y \cdot \{x, z\}$  for all  $x, y, z$  in the Poisson algebra.

**Exercise 7**

Prove that both methods of calculating  $\dot{f}$  yield the same result.

**Solution.**

$$\begin{aligned} \frac{df}{dt} &= \dot{g}h + g\dot{h} \\ &= \{g, H\}h + g\{h, H\} \\ &= -(\{H, g\}h + g\{H, h\}) \\ &= -\{H, gh\} \\ &= \{gh, H\} = \{f, H\} \end{aligned}$$

**Exercise 8**

Use the Jacobi identity to prove that

$$\frac{d}{dt}\{f, g\} = \{\dot{f}, g\} + \{f, \dot{g}\}$$

**Solution.**

$$\begin{aligned}
 \frac{d}{dt}\{f, g\} &= \{\{f, g\}, H\} \\
 &= -\{H, \{f, g\}\} \\
 &= \{f, \{g, H\}\} + \{g, \{H, f\}\} && \text{(Jacobi identity)} \\
 &= \{f, \dot{g}\} + \{g, -\dot{f}\} \\
 &= \{\dot{f}, g\} + \{f, \dot{g}\}
 \end{aligned}$$

### Exercise 9

Show that if  $H$  is a polynomial in the positions and momenta with arbitrary (and possibly time-dependent) coefficients, it is true that  $\frac{dH}{dt} = \frac{\partial H}{\partial t}$ .

**Solution.**

$$\begin{aligned}
 \frac{d}{dt}H(x, p, t) &= \frac{\partial H}{\partial x} \frac{dx}{dt} + \frac{\partial H}{\partial p} \frac{dp}{dt} + \frac{\partial H}{\partial t} \\
 &= \frac{\partial H}{\partial x} \dot{x} + \frac{\partial H}{\partial p} \dot{p} + \frac{\partial H}{\partial t} \\
 &= \frac{\partial H}{\partial x} \frac{\partial H}{\partial p} - \frac{\partial H}{\partial p} \frac{\partial H}{\partial x} + \frac{\partial H}{\partial t} \\
 &= \frac{\partial H}{\partial t}
 \end{aligned}$$

### Exercise 10

Show that the total momentum is conserved.

**Solution.** To show  $p_i^{(1)} + p_i^{(2)}$  is conserved for all  $i$  we will show it's time derivative is 0 by taking it's Poisson bracket with the Hamiltonian.

$$\begin{aligned}
 \frac{d}{dt}(p_i^{(1)} + p_i^{(2)}) &= \{p_i^{(1)} + p_i^{(2)}, H\} \\
 &= \left\{ p_i^{(1)} + p_i^{(2)}, \frac{k}{2} \sum_j (x_j^{(1)} - x_j^{(2)})^2 \right\} \\
 &= \frac{k}{2} \left\{ p_i^{(1)} + p_i^{(2)}, (x_i^{(1)} - x_i^{(2)})^2 \right\} \\
 &= \frac{k}{2} \left[ \left\{ p_i^{(1)}, x_i^{(1)2} \right\} - 2 \left\{ p_i^{(1)}, x_i^{(1)} x_i^{(2)} \right\} + \left\{ p_i^{(2)}, x_i^{(2)2} \right\} - 2 \left\{ p_i^{(2)}, x_i^{(1)} x_i^{(2)} \right\} \right] \\
 &= \frac{k}{2} \left[ -2x_i^{(1)} + 2x_i^{(2)} - 2x_i^{(2)} + 2x_i^{(1)} \right] \\
 &= 0
 \end{aligned}$$

With this, and the fact that  $H$  has no explicit time dependence, we can conclude that momentum is conserved.

**Exercise 11**

Use the derivative definition of the Poisson bracket to evaluate  $\{x^8 p^6, x^3 p^4\}$ .

**Solution.**

$$\begin{aligned}\{x^8 p^6, x^3 p^4\} &= \frac{\partial}{\partial x}(x^8 p^6) \frac{\partial}{\partial p}(x^3 p^4) - \frac{\partial}{\partial p}(x^8 p^6) \frac{\partial}{\partial x}(x^3 p^4) \\ &= (8x^7 p^6)(4x^3 p^3) - (6x^8 p^5)(3x^2 p^4) \\ &= 32x^{10} p^9 - 18x^{10} p^9 \\ &= 14x^{10} p^9\end{aligned}$$

**Exercise 12**

Show that the derivative definition of the Poisson bracket is indeed a representation of the Poisson bracket defined by the axioms.

**Solution.** First, antisymmetry.

$$\begin{aligned}\{f, g\} &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial g}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial g}{\partial x_i^{(r)}} \right) \\ &= - \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial g}{\partial x_i^{(r)}} \frac{\partial f}{\partial p_i^{(r)}} - \frac{\partial g}{\partial p_i^{(r)}} \frac{\partial f}{\partial x_i^{(r)}} \right) \\ &= -\{g, f\}\end{aligned}$$

Second, linearity.

$$\begin{aligned}\{cf, g\} &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial(cf)}{\partial x_i^{(r)}} \frac{\partial g}{\partial p_i^{(r)}} - \frac{\partial(cf)}{\partial p_i^{(r)}} \frac{\partial g}{\partial x_i^{(r)}} \right) \\ &= c \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial g}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial g}{\partial x_i^{(r)}} \right) \\ &= c\{f, g\}\end{aligned}$$

Third, the addition rule.

$$\begin{aligned}\{f, g+h\} &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial(g+h)}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial(g+h)}{\partial x_i^{(r)}} \right) \\ &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial g}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial g}{\partial x_i^{(r)}} + \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial h}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial h}{\partial x_i^{(r)}} \right) \\ &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial g}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial g}{\partial x_i^{(r)}} \right) + \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial h}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial h}{\partial x_i^{(r)}} \right) \\ &= \{f, g\} + \{f, h\}\end{aligned}$$

Fourth, the product rule.

$$\begin{aligned}
 \{f, gh\} &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial(gh)}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial(gh)}{\partial x_i^{(r)}} \right) \\
 &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \left[ \frac{\partial g}{\partial p_i^{(r)}} h + g \frac{\partial h}{\partial p_i^{(r)}} \right] - \frac{\partial f}{\partial p_i^{(r)}} \left[ \frac{\partial g}{\partial x_i^{(r)}} h + g \frac{\partial h}{\partial x_i^{(r)}} \right] \right) \\
 &= h \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial g}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial g}{\partial x_i^{(r)}} \right) \\
 &\quad + g \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial h}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial h}{\partial x_i^{(r)}} \right) \\
 &= \{f, g\}h + g\{f, h\}
 \end{aligned}$$

Fifth, sixth, and seventh: the rules for positions and momenta.

$$\begin{aligned}
 \{x_i^{(r)}, p_j^{(s)}\} &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial x_i^{(r)}}{\partial x_i^{(r)}} \frac{\partial p_j^{(s)}}{\partial p_i^{(r)}} - \frac{\partial x_i^{(r)}}{\partial p_i^{(r)}} \frac{\partial p_j^{(s)}}{\partial x_i^{(r)}} \right) \\
 &= \sum_{r=1}^n \sum_{i=1}^3 \delta_{i,j} \delta_{s,r} \\
 &= \delta_{i,j} \delta_{s,r}
 \end{aligned}$$

$$\begin{aligned}
 \{x_i^{(r)}, x_j^{(s)}\} &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial x_i^{(r)}}{\partial x_i^{(r)}} \frac{\partial x_j^{(s)}}{\partial p_i^{(r)}} - \frac{\partial x_i^{(r)}}{\partial p_i^{(r)}} \frac{\partial x_j^{(s)}}{\partial x_i^{(r)}} \right) \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 \{p_i^{(r)}, p_j^{(s)}\} &= \sum_{r=1}^n \sum_{i=1}^3 \left( \frac{\partial p_i^{(r)}}{\partial x_i^{(r)}} \frac{\partial p_j^{(s)}}{\partial p_i^{(r)}} - \frac{\partial p_i^{(r)}}{\partial p_i^{(r)}} \frac{\partial p_j^{(s)}}{\partial x_i^{(r)}} \right) \\
 &= 0
 \end{aligned}$$

### Exercise 13

Find the representation of the Hamilton equations

**Solution.** We start with eq. 2.21 and eq. 2.22 and then use those results for eq. 2.19.

$$\begin{aligned}
 \frac{d}{dt}x_i^{(r)} &= \{x_i^{(r)}, H\} \\
 &= \sum_{s,j} \frac{\partial x_i^{(r)}}{\partial x_j^{(s)}} \frac{\partial H}{\partial p_j^{(s)}} - \cancel{\frac{\partial x_i^{(r)}}{\partial p_j^{(s)}} \frac{\partial H}{\partial x_j^{(s)}}} \\
 &= \sum_{s,j} \delta_{i,j} \delta_{r,s} \frac{\partial H}{\partial p_j^{(s)}} \\
 \dot{x}_i^{(r)} &= \frac{\partial H}{\partial p_i^{(r)}}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d}{dt}p_i^{(r)} &= \{p_i^{(r)}, H\} \\
 &= \sum_{s,j} \cancel{\frac{\partial p_i^{(r)}}{\partial x_j^{(s)}} \frac{\partial H}{\partial p_j^{(s)}}} - \frac{\partial p_i^{(r)}}{\partial p_j^{(s)}} \frac{\partial H}{\partial x_j^{(s)}} \\
 &= - \sum_{s,j} \delta_{i,j} \delta_{r,s} \frac{\partial H}{\partial x_j^{(s)}} \\
 \dot{p}_i^{(r)} &= - \frac{\partial H}{\partial x_i^{(r)}}
 \end{aligned}$$

$$\begin{aligned}
 \frac{df}{dt} &= \{f, H\} + \frac{\partial f}{\partial t} \\
 &= \sum_{r,i} \frac{\partial f}{\partial x_i^{(r)}} \frac{\partial H}{\partial p_i^{(r)}} - \frac{\partial f}{\partial p_i^{(r)}} \frac{\partial H}{\partial x_i^{(r)}} + \frac{\partial f}{\partial t} \\
 &= \sum_{r,i} \frac{\partial f}{\partial x_i^{(r)}} \dot{x}_i^{(r)} + \frac{\partial f}{\partial p_i^{(r)}} \dot{p}_i^{(r)} + \frac{\partial f}{\partial t}
 \end{aligned}$$

Which is exactly the chain rule for a function  $f(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)}, \mathbf{p}^{(1)}, \dots, \mathbf{p}^{(n)}, t)$ .